

1 **Perfluoroalkyl Acids are Common Environmental Pollutants**
2 **That Potently Stimulate Aldosterone Biosynthesis**

3
4 Brasilina Caroccia¹, Teresa Maria Seccia¹, Giorgia Pallafacchina^{2,3}, Maria Piazza¹, Ilaria
5 Caputo¹, Stefania Zamberlan¹, Rosario Rizzuto² and Gian Paolo Rossi¹
6

7
8 ¹Internal Emergency Medicine Unit, Department of Medicine-DIMED University of Padua, Specialized Center
9 for Blood Pressure Disorders-Regione Veneto, Padua, Italy

10 ²Department of Biomedical Sciences-DSB, University of Padua, Padua, Italy


11 ³Italian National Research Council (CNR), Neuroscience Institute, Padua, Italy.
12

13 **Running title:** Perfluoroalkyl Substances and Aldosterone
14
15
16
17
18
19
20

21 **Corresponding author:**

22 Professor Gian Paolo Rossi, MD, FACC, FAHA.
23 Internal Emergency Medicine Unit, Department of Medicine-DIMED
24 University of Padua,
25 & Specialized Center for Blood Pressure Disorders-Regione Veneto,
26 University Hospital
27 Via Giustiniani, 2
28 35126 Padova, Italy
29 Phone: +39-049-821-2279 or 7821
30 Fax: +39-049-8217873
31 E-mail: gianpaolo.rossi@unipd.it
32
33
34
35
36
37
38
39
40

41
42
43



44 **Supplemental Figure 1. Adrenocortical cells viability is not affected by perfluoroalkyl**
45 **substances.**

46 HAC15 cell viability was evaluated by MTT assay at 3 different time points (24, 48 or 72 hours)
47 after treatment with two different concentration, 1 μM (in light grey) or 10 μM (in dark grey),
48 of PFOA (Panel A), PFOS (Panel B) and PFAS (PFOA + PFOS, Panel C). Cell viability was >95% in
49 all the conditions analysed. Data are reported as a percentage of the treated cells on the
50 untreated controls. One-way ANOVA with Dunnett post hoc test was performed and no
51 significant statistical difference was observed.